



⑪ Publication number:

0 098 110
B1

⑫ **EUROPEAN PATENT SPECIFICATION**

④⑤ Date of publication of patent specification: **18.10.89**

②① Application number: **83303636.1**

⑦② Date of filing: **23.06.83**

⑤① Int. Cl.⁴: **A 61 K 47/00, A 61 K 37/02,**
A 61 K 37/24, A 61 K 37/47,
A 61 K 37/48, A 61 K 37/64,
C 12 N 11/08, A 61 K 45/02

⑤④ Long-acting composition.

③⑥ Priority: **24.06.82 JP 108851/82**
27.09.82 JP 169160/82

④③ Date of publication of application:
11.01.84 Bulletin 84/02

④⑤ Publication of the grant of the patent:
18.10.89 Bulletin 89/42

④④ Designated Contracting States:
CH DE FR GB IT LI

⑤③ References cited:
EP-A-0 018 609
WO-A-83/00288
DE-A-1 802 465
FR-A-2 515 684

CHEMICAL ABSTRACTS, vol. 91, no. 7, 13th
August 1979, pages 281-282, no. 51928v,
Columbus, Ohio, US; K.V. SAVOCA et al.:
"Preparation of a non-immunogenic arginase
by the covalent attachment of polyethylene
glycol", & BIOCHIM. BIOPHYS. ACTA. 1979,
578(1), 47-53

The file contains technical information
submitted after the application was filed and
not included in this specification

⑦③ Proprietor: **NIHON CHEMICAL RESEARCH**
KABUSHIKI KAISHA also known as **JAPAN**
CHEMICAL RESEARCH CO., LTD
4-20, Mikagehommachi 3-chome Higashinada-
ku
Kobe Hyogo 658 (JP)

⑦② Inventor: **Hiratani, Hajime**
705-3, Tottori Hannan-cho
Sennan-gun Osaka 559-02 (JP)

⑦④ Representative: **Burford, Anthony Frederick**
et al
W.H. Beck, Greener & Co. 7 Stone Buildings
Lincoln's Inn
London WC2A 3SZ (GB)

⑤③ References cited:
CHEMICAL ABSTRACTS, vol. 87, no. 7, 15th
August 1977, page 161, no. 49460k, Columbus,
Ohio, US; A. ABUCHOWSKI et al.: "Effect of
covalent attachment of polyethylene glycol on
immunogenicity and circulating life of bovine
liver catalase", & J. BIOL. CHEM. 1977, 252(11),
3582-6

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may
give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall
be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been
paid. (Art. 99(1) European patent convention).

Courier Press, Leamington Spa, England.

EP 0 098 110 B1

Description

This invention relates to a long-active composition which is effective in the human body for a prolonged period of time. More particularly, this invention relates to a long-active composition wherein a physiologically active polypeptide or glycoprotein of human origin is coupled to a polyoxyethylene-polyoxypropylene copolymer.

Physiologically active substances, such as a variety of hormones and enzymes, when administered to living bodies, generally undergo decomposition by various proteases in said living bodies in a short time or immediate inhibition by various inhibitors, so that their effects can last only for a short period of time.

It has been disclosed by Abuchowski et al of Department of Biochemistry, Rutgers, State University, New Brunswick, N.J., U.S.A. (CA 87:49460k and CA 91:51928v) that bovine liver catalase and arginase can be stabilized by covalent attachment to methoxypolyethylene glycol (MW1900 or MW5000) with cyanuric chloride as the coupling agent.

The present inventors conducted intensive research in search of the possibility of rendering their physiological activity lasting in living bodies so that their effects can be ensured to an increased extent and their doses can be reduced. As a result, it has been found that the above objects can be accomplished by coupling physiologically active substances to polyoxyethylene-polyoxypropylene copolymers (hereinafter briefly referred to as "copolymers").

The present invention thus provides long-acting compositions comprising human-derived physiologically active polypeptides or glycoproteins coupled to polyoxyethylene-polyoxypropylene copolymers.

In another aspect thereof, the present invention provides a method of rendering the above-mentioned polypeptides or glycoproteins long-acting by coupling them to the above-mentioned copolymers.

Polyoxyethylene-polyoxypropylene copolymers are available on the market. About 33 species are in main use; they have a molecular weight of 1,000 to 14,000 with varied polyoxyethylene-to-polyoxypropylene ratios, e.g. 4:16, 196:67 or 256:54. However, products obtained by coupling physiologically active substances to those copolymers which have a molecular weight exceeding 10,000 have undesirably decreased activity due to their active groups being wrapped up in or concealed by the copolymer molecules. Such products are undesirable also because they remain in living bodies for an excessively long period and possibly produce side effects.

Accordingly, in accordance with the present invention, those copolymers that have a molecular weight of 1,000 to 10,000 are used. The copolymers have hydroxyl groups at both the ends thereof. The hydrogen atom of one of the hydroxyl groups may be substituted by an alkyl or acyl group. Preferred examples of the alkyl group

are methyl and ethyl. Acetyl and propionyl are examples of the acyl group. Substitution by these groups can be performed by the known methods.

In practicing the present invention, a polypeptide or glycoprotein of human origin, e.g. extracted from human urine, placenta or blood components are derived from human blood components or produced by tissue culture of human cells, is used.

Examples of the polypeptide or glycoprotein are human menopausal gonadotropin (HMG), human growth hormone (HGH), epidermal growth factor (EGF), nerve growth factor (NGF), colony formation stimulating factor (CSF), urokinase (UK), plasminogen (PLG), kallikrein, interferon α , interferon β , interferon γ , interleukin 1, interleukin 2, urinary trypsin inhibitor, urinary thiol protease inhibitor, placental arylsulfatase, urinary lysozyme and urinary asparaginase.

In accordance with the present invention, the above-mentioned copolymers are coupled to such physiologically active polypeptides or glycoproteins. The coupling is effected by means of a coupling agent capable of linking a terminal hydroxyl group of the copolymer with the amino group of the active substance. The coupling agent contains at least one functional group capable of reacting with the hydroxyl group and at least one functional group capable of reacting with the amino group, and includes 2,4,6-trichloro-s-triazine, dibromosuccinic anhydride and maleic anhydride.

Thus, for instance, the copolymer is reacted with 2,4,6-trichloro-s-triazine in the presence of an alkali, and the resulting reactive copolymer derivative is reacted with the above-mentioned active substance, whereby the active substance is coupled to at least one copolymer molecule through the N-terminal primary amino group and/or the ϵ -amino group or groups in a lysine residue or residues.

The above coupling reaction can be performed by known methods, the reaction conditions being selected based on the reactivity each of the terminal hydroxyl group of the copolymer, the amino group of the active substance, and the coupling agent.

The compositions according to the present invention can produce the effects of the active substances in living bodies for a significantly prolonged period of time; the duration of activity can be prolonged 10 to 20 times, for instance.

Moreover, the copolymer-physiologically active substance coupling products are rather inert to proteases in living bodies and furthermore are not susceptible to various inhibitors in the blood, and therefore said products show marked improvements in development and persistence of activity.

The compositions according to the present invention can be administered orally or parenterally depending on the kind of physiologically active substances. Parenteral administration may be conducted in the manner of intravenous, intramuscular or subcutaneous injection.

The dose of the compositions is proportional to the dose known for the physiologically active substances concerned. However, the compositions according to the present invention tend to have a somewhat decreased activity per unit weight as compared with the starting active substances and therefore it is desirable to increase the single dose thereof accordingly. Nevertheless, since they are markedly improved in the duration of effects, as mentioned above, the compositions may be administered every several days or at longer intervals where the corresponding active substances themselves should be administered daily.

The following examples illustrate the present invention in more detail.

Example 1

2,4,6-Trichloro-s-triazine (cyanuric chloride) (5.5 g, 30 millimoles) was added to 400 ml of anhydrous benzene containing 10 g of anhydrous sodium carbonate, followed by addition of 50 g (10 millimoles) of methoxy-polyoxyethylene-polyoxypropylene glycol [average molecular weight 5,000, monomethylation product derived from Pluronic F-38 (Asahi Denka Kogyo K.K.), E.O.:P.O.:E.O.=46:16:46]. The mixture was stirred at room temperature overnight.

The insoluble matter was filtered off, 5 volumes of petroleum ether was added to the filtrate and the resulting precipitate, activated copolymer, was collected by filtration. Two more redissolution-precipitation procedures with benzene and petroleum ether gave 51.5 g of the desired activated copolymer.

Purified urokinase (three million units) was dissolved in 30 ml of 0.1 M phosphate buffer (pH 7.0 at 4°C. To the solution was added 600 mg of the above activated copolymer, and the mixture was stirred at 4°C with stirring for 3 hours.

Then, the reaction was terminated by adjusting the pH to not more than 5.0. The unreacted activated copolymer was removed by gel filtration using Sephadex G-100 equilibrated with 0.1 M phosphate buffer (pH 5.0).

The thus-obtained modified urokinase had an average molecular weight of 150,000 and retained 40% (fibrin plate method) or 70% (fluorogenic substrate method) of the original activity.

The half lives in the blood of the unmodified urokinase and the modified urokinase as determined in rabbits were 5 minutes and 120 minutes, respectively, the half life ratio thus being as high as 24. The half life determination was conducted in the following manner.

The unmodified or modified urokinase (50,000 units/kg) was administered to rabbits weighing about 2.0 kg into the auricular vein on the side opposite to the ear where blood sampling was conducted at timed intervals.

Blood sampling was performed by connecting a syringe with an indwelling needle in the auricular vein. Ten to thirty minutes before urokinase administration, heparin sodium was intravenously administered in a dose of 1,000 units/kg.

Before administration of the sample, immediately after administration, and 2 minutes, 5 minutes, 10 minutes, 20 minutes, 30 minutes, 40 minutes, 60 minutes, 120 minutes and 240 minutes after administration, 2 ml of blood was collected from each animal and immediately centrifuged (3,000 rpm, 5 minutes), and the plasma was assayed for urokinase activity. The above-mentioned results were obtained in that manner.

The fibrin plate method used in this example for urokinase activity determination was performed as described by P. L. Walton in Clin. Chim. Acta, 13 (5), 680-684 (1966).

The fluorogenic substrate method was performed as described by T. Morita et al. in J. Biochem., 82, 1495 (1977).

Example 2

100,000 units of human urinary kallikrein purified almost to absolute purity was dissolved in 50 ml of 0.1 M phosphate buffer (pH 7.0) at 4°C. Ethoxy-polyoxyethylene-polyoxypropylene glycol (average molecular weight 3,400, E.O.:P.O.:E.O.=19:30:19, Asahi Denka Kogyo's Pluronic P-65) was activated in the same manner as in Example 1. The activated copolymer (0.7 g) was added to the above solution, and the mixture was stirred at 4°C for 3 hours.

Then, the reaction was terminated by adjusting the pH to not more than 5.0, and the unreacted activated copolymer was removed by dialyzing the reaction mixture at 4° overnight against 0.1 M phosphate buffer pH 5.0.

The thus-obtained modified kallikrein had an average molecular weight of 100,000 and retained 50% (hypotension method using dogs) or 80% (fluorogenic substrate method using Pro-Phe-Arg-MCA) of the original activity.

The half lives of the unmodified kallikrein and the modified kallikrein were determined in rabbits by collecting blood samples in the same manner as in Example 1. They were 7 minutes and 110 minutes, respectively, and the ratio therebetween amounted to 15.

The hypotension method using dogs as used in this example for the assay of kallikrein was performed as described in J. Biochem., 58, 201 (1965).

The fluorogenic substrate method was performed as described in J. Biochem., 82, 1495 (1977).

Example 3

100 million units of human leukocyte interferon (specific activity 2×10^7 units/mg protein) was dissolved in 22 ml of 0.1 M phosphate buffer (pH 7.0) at 4°C, then 220 mg of the same activated copolymer as used in Example 1 was added, and the mixture was stirred at 4°C for 3 hours. Then, the reaction was terminated by adjusting the pH to not more than 5.0, and the unreacted activated copolymer was removed by subjecting the reaction mixture to dialysis against 0.1 M phosphate buffer (pH 5.0) at 4°C overnight. The thus-obtained modified interferon retained 40% of

the activity before modification. The cells used in the activity determination were FL-cells [human amniotic cells (Fogh and Lund strain)]. VSV (vesicular stomatitis virus) was used as the challenging virus. The CPE (cytopathogenic effect) manifested by the microplate method was judged by the dye uptake method [N. B. Finter, J. General Virology, 5, 419 (1969)] using phenol red. The half lives in rabbit blood of the unmodified interferon α and the modified interferon α as determined with blood samples collected in the same manner as in Example 1 were 5 minutes and 80 minutes, respectively, revealing a 16 times prolonged half life of the modified one.

Claims

1. A long-acting composition comprising a physiologically active polypeptide or glycoprotein coupled to a polymer characterised in that the polypeptide or glycoprotein is coupled to a polyoxyethylene-polyoxypropylene copolymer, having a molecular weight of 1,000 to 10,000 and having at both ends thereof hydroxyl groups one of which may be substituted by an alkyl or acyl group, by means of a coupling agent linking a terminal hydroxyl group of said copolymer with an amino group of said active substance.

2. A composition as claimed in Claim 1, wherein the physiologically active polypeptide or glycoprotein is of human origin.

3. A composition as claimed in Claim 2, wherein said polypeptide or glycoprotein is human menopausal gonadotropin, human growth hormone, epidermal growth factor, nerve growth factor, colony formation stimulating factor, urokinase, plasminogen (PLG), kallikrein, interferon α , interferon β , interferon γ , interleukin-1, interleukin 2, urinary trypsin inhibitor, urinary thiol protease inhibitor, placental arylsulfatase, urinary lysozyme or urinary asparaginase.

4. A composition as claimed in Claim 2, wherein the physiologically active polypeptide or glycoprotein is urokinase, kallikrein or leukocyte interferon.

5. A composition as claimed in any one of the preceding claims, wherein the copolymer has a polyoxyethylene-polyoxypropylene ratio from 4:16 to 256:54.

6. A composition as claimed in any one of the preceding claims, wherein copolymer is a methoxypolyoxyethylene-polyoxypropylene copolymer having E.O.:P.O.:E.O. ratio of 46:16:46 and has average molecular weight of 5,000.

7. A composition as claimed in any one of the preceding claims, wherein the copolymer is an ethoxypolyoxyethylene-polyoxypropylene glycol having average molecular weight of 3,400 and E.O.:P.O.:E.O. ratio of 19:30:19.

8. A composition as claimed in any one of the preceding claims, wherein the coupling agent contains at least one functional group capable of reacting with a hydroxyl group and at least one functional group capable of reacting with an amino group.

9. A composition as claimed in Claim 8, wherein the coupling agent is 2,4,6-trichloro-s-triazine, dibromosuccinic anhydride or maleic anhydride.

Patentansprüche

1. Lang wirkendes Mittel, umfassend ein an ein Polymer gekuppeltes physiologisch aktives Polypeptid oder Glycoprotein, dadurch gekennzeichnet, daß das Polypeptid oder Glycoprotein mit Hilfe eines Kupplungsmittels an ein Polyoxyethylen-polyoxypropylen-copolymer gebunden ist, das ein Molekulargewicht von 1 000 bis 10 000 hat und an beiden Enden Hydroxylgruppen aufweist, wovon eine durch eine Alkyl- oder Acylgruppe substituiert sein kann, wobei das Kupplungsmittel eine terminale Hydroxylgruppe des Copolymers mit einer Aminogruppe der aktiven Substanz verbindet.

2. Mittel nach Anspruch 1, worin das physiologisch aktive Polypeptid oder Glycoprotein menschlichen Ursprungs ist.

3. Mittel nach Anspruch 1, wobei es sich bei dem Polypeptid oder Glycoprotein um menopausales Humangonadotropin, menschliches Wachstumshormon, den epidermalen Wachstumsfaktor, der Nervenwachstumsfaktor, den die Koloniebildung anregenden Faktor, Urokinase, Plasminogen (PLG), Kallikrein, Interferon α , Interferon β , Interferon γ , Interleukin 1, Interleukin 2, den Harntrypsin-Inhibitor, den Harnthiolprotease-Inhibitor, placentare Arylsulfatase, Harnlysozym oder Harnasparaginase handelt.

4. Mittel nach Anspruch 2, wobei es sich bei dem physiologisch aktiven Polypeptid oder Glycoprotein um Urokinase, Kallikrein oder Leukocyten-Interferon handelt.

5. Mittel nach einem der vorhergehenden Ansprüche, wobei das Polyoxyethylen-Polyoxypropylen-Verhältnis des Copolymers 4:16 bis 256:54 ist.

6. Mittel nach einem der vorhergehenden Ansprüche, wobei es sich bei dem Copolymer um ein Methoxypolyoxyethylen-polyoxypropylen-copolymer mit einem E.O.:P.O.:E.O.-Verhältnis von 46:16:46 und einem durchschnittlichen Molekulargewicht von 5 000 handelt.

7. Mittel nach einem der vorhergehenden Ansprüche, wobei es sich bei dem Copolymer um Ethoxypolyoxyethylen-polyoxypropylen-glycol mit einem durchschnittlichen Molekulargewicht von 3 400 und einem E.O.:P.O.:E.O.-Verhältnis von 19:30:19 handelt.

8. Mittel nach einem der vorhergehenden Ansprüche, worin das Kupplungsmittel mindestens eine zur Reaktion mit einer Hydroxylgruppe befähigte funktionelle Gruppe und mindestens eine zur Reaktion mit einer Aminogruppe befähigte funktionelle Gruppe enthält.

9. Mittel nach Anspruch 8, wobei es sich bei dem Kupplungsmittel um 2,4,6-Trichlor-s-triazin, Dibrombernsteinsäureanhydrid oder Maleinsäureanhydrid handelt.

Revendications

1. Composition à longue activité, comprenant un polypeptide ou une glycoprotéine physiologiquement actif, couplé à un polymère, caractérisée en ce que le polypeptide ou la glycoprotéine est couplé à un copolymère polyoxyéthylène-polyoxypropylène ayant un poids moléculaire de 1.000 à 10.000 et présentant à ses deux extrémités, des groupes hydroxyle, dont l'un peut être substitué par un groupe alkyle ou acyle, au moyen d'un agent de couplage liant un groupe hydroxyle terminal dudit copolymère à un groupe amino de ladite substance active.

2. Composition selon la revendication 1, caractérisée en ce que le polypeptide ou la glycoprotéine physiologiquement actif est d'origine humaine.

3. Composition selon la revendication 2, caractérisée en ce que ledit polypeptide ou ladite glycoprotéine est la gonadotropine ménopausique humaine, l'hormone de croissance humaine, un facteur de croissance épidermique, un facteur de croissance nerveux, un facteur stimulant la formation de colonies, l'urokinase, le plasminogène (PLG), la kallikréine, l'interféron α , l'interféron β , l'interféron γ , l'interleucine 1, l'interleucine 2, l'inhibiteur de trypsine urinaire, l'inhibiteur de thiol protéase urinaire, une arylsulfatase placentaire, une lysozyme urinaire ou une asparaginase urinaire.

4. Composition selon la revendication 2, caractérisée en ce que le polypeptide ou la glycoprotéine physiologiquement actif est l'urokinase, la kallikréine ou un interféron leucocyte.

5. Composition selon l'une des revendications précédentes, caractérisée en ce que le copolymère présente un rapport polyoxyéthylène-polyoxypropylène de 4:16 à 256:54.

6. Composition selon l'une des revendications précédentes, caractérisée en ce que le copolymère est un copolymère méthoxypolyoxyéthylène-polyoxypropylène présentant un rapport E.O.:P.O.:E.O. de 46:16:46 et un poids moléculaire moyen de 5.000.

7. Composition selon l'une des revendications précédentes, caractérisée en ce que le copolymère est un éthoxypolyoxyéthylène-polyoxypropylène glycol ayant un poids moléculaire moyen de 3.400 et un rapport E.O.:P.O.:E.O. de 19:30:19.

8. Composition selon l'une des revendications précédentes, caractérisée en ce que l'agent de couplage contient au moins un groupe fonctionnel capable de réagir avec un groupe hydroxyle et au moins un groupe fonctionnel capable de réagir avec un groupe amino.

9. Composition selon la revendication 8, caractérisée en ce que l'agent de couplage est la 2,4,6-trichloro-s-triazine, l'anhydride dibromosuccinique ou l'anhydride maléique.

35

40

45

50

55

60

65

5